

Lymphoid involvement of cerebrospinal fluid in Lyme disease

Neuroborréliose : infiltration lymphoïde pléomorphe du LCR

Floriane Lefeuvre

Lucile Baseggio

Groupement hospitalier Lyon-Sud,
Hospices civils de Lyon, Laboratoire
d'hématologie biologique, France

Article received August 22, 2019,
accepted December 04, 2019

After typical rash (erythema migrans), neurological involvement is the most common clinical manifestation in Lyme disease. The cerebrospinal fluid (CSF) analysis may help in any diagnostic doubt characterized by CSF pleocytosis (lymphoïd and plasma cells) and *borrelia*-specific antibodies produced intathecally.

A 62-year-old man presented with a back pain radiating to right thigh, severe asthenia and no fever. Serum C protein and erythrocyte sedimentation rate were normal. Two months later given tetraparesis manifestation, a MR scan of spine has been realized and displayed high-density signal extending between T11 to L1. The MR scan of brain was normal. Infectious mononucleosis screen, syphilis, HIV, hepatitis B and C serology were negative. Cerebrospinal fluid (CSF) revealed hypercellular specimen (white blood cells=630/ μ L, red blood cells=10/ μ L). The protein concentration was 2.1 g/L and glucose 3.2 mmol/L (normal paired plasma glucose). The cytological analysis of CSF (cytocentrifuge preparation, magnification x60, May-Grünwald Giemsa staining, *figure 1 A-C*) showed polymorphous lymphoïd population including small lymphocytes, granular lymphocytes, large immunoblasts (arrow) and plasma cells

(asterisk). Flow cytometry (FCM) immunophenotyping of CSF (*figure 1 D*) identified mainly small T-cells (84%) associated to polytypic B-cells (10%) and polytypic plasma cells (4%). Morphology associated to immunologic profile argue to CSF involvement by activated lymphoïd population. The patient only remembered untreated skin lesions on abdomen 10 weeks before. The *Borrelia* enzyme immunoassay result was found to be reactive and confirmed with positive immunoblot to IgG and IgM. The CSF also contained antibodies to *Borrelia* and the *Borrelia*-specific CSF to serum IgG antibody index was at 8.1 (normal range <1.3). The patient was therefore diagnosed with Lyme neuroborreliosis and treated with intravenous ceftriaxone, which resulted in an almost complete recovery.

In conclusion, this case illustrates CNS involvement in neuroborreliosis that will develop in less of 5% of untreated erythema migrans, and usually 4-6 weeks after tick exposure [1]. Although the analysis of CSF is not recommended in all patients, in any diagnostic doubt it will be usefulness with characteristically CSF pleocytosis (lymphoïd and plasma cells) and *borrelia*-specific antibodies produced intathecally [2].

Correspondance : L. Baseggio
<lucile.baseggio@chu-lyon.fr>




Figure 1. Cerebrospinal fluid (CSF) (cytocentrifuge preparation, magnification x60, May-Grünwald Giemsa staining, **A-C**): small lymphoid cells (scanty cytoplasm and regular nuclear), granular lymphocytes (abundant and pale cytoplasm with azurophilic granules), immunoblasts (large cells with abundant and basophilic cytoplasm; arrow) and plasma cells (eccentric nuclei with clumped chromatin and basophilic cytoplasm; asterisk). Immunophenotype by multiparameter flow cytometry (FCM) was realized on CSF (**D**). Data were acquired on a BD FACSCanto II cytometer and analyzed with BD FACS DIVA version 8.0.1 software (*Becton Dickinson, San Jose, CA, USA). A first panel of 10 antibodies in one tube was used to identify T- and B-cells : CD3 (clone SK7, 15-18*, APC-H7), CD5 (clone L17F12*, PerCP Cy5.5), CD4 (clone RPA-T4*, V450), CD8 (RPA-T8*, FITC), CD45 (clone HI30*, V500), CD20 (clone L27-9*, V450), CD19 (clone J3-119**, PE Cy7), CD10 (clone HI10a3*, APC), CD56 (clone NCAM 16,2*, PE) and kappa/lambda (**, FITC/PE) (** Beckman-Coulter, Hialeah, FL, USA, **Dako, Carpinteria, CA). A second panel was used to identify plasma cells (including CD38 [clone HB7*, V450], CD138 [Syndecan-1**, PC5]).

Letter

Conflict of interest: none of the authors has any conflict of interest to disclose.

2. NICE, 2018. Lyme disease overview. <https://pathways.nice.org.uk/pathways/lyme-disease>.

Reference

1. Hansen K, Lebech AM. The clinical and epidemiological profile of Lyme neuroborreliosis in Denmark 1985-1990, (1992) A prospective study of 187 patients with *Borrelia burgdorferi* specific intrathecal antibody production. *Brain* 1992; 115: 399-423.